

WEST Search History

DATE: Thursday, May 29, 2003

<u>Set Name</u> side by side	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u> result set
<i>DB=USPT,PGPB; PLUR=YES; OP=ADJ</i>			
L13	L12 and stress	24	L13
L12	L2 and transgenic	91	L12
L11	L10 and stress	25	L11
L10	L2 and antisense	83	L10
L9	L2 and turgidity	2	L9
L8	l2 and abscisic acid	6	L8
L7	l2 and abscisic acid	0	L7
L6	L2 and diffusion resistance	1	L6
L5	l2 and transpiration	1	L5
L4	l2 and drought	9	L4
L3	L2 and stomat\$	9	L3
L2	L1 and plant	429	L2
L1	p1d or phospholipase d	5612	L1

END OF SEARCH HISTORY

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:sssptal649axm

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS	1		Web Page URLs for STN Seminar Schedule - N. America
NEWS	2	Apr 08	"Ask CAS" for self-help around the clock
NEWS	3	Jun 03	New e-mail delivery for search results now available
NEWS	4	Aug 08	PHARMAMarketLetter(PHARMAML) - new on STN
NEWS	5	Aug 19	Aquatic Toxicity Information Retrieval (AQUIRE) now available on STN
NEWS	6	Aug 26	Sequence searching in REGISTRY enhanced
NEWS	7	Sep 03	JAPIO has been reloaded and enhanced
NEWS	8	Sep 16	Experimental properties added to the REGISTRY file
NEWS	9	Sep 16	CA Section Thesaurus available in CAPLUS and CA
NEWS	10	Oct 01	CASREACT Enriched with Reactions from 1907 to 1985
NEWS	11	Oct 24	BEILSTEIN adds new search fields
NEWS	12	Oct 24	Nutraceuticals International (NUTRACEUT) now available on STN
NEWS	13	Nov 18	DKILIT has been renamed APOLLIT
NEWS	14	Nov 25	More calculated properties added to REGISTRY
NEWS	15	Dec 04	CSA files on STN
NEWS	16	Dec 17	PCTFULL now covers WP/PCT Applications from 1978 to date
NEWS	17	Dec 17	TOXCENTER enhanced with additional content
NEWS	18	Dec 17	Adis Clinical Trials Insight now available on STN
NEWS	19	Jan 29	Simultaneous left and right truncation added to COMPENDEX, ENERGY, INSPEC
NEWS	20	Feb 13	CANCERLIT is no longer being updated
NEWS	21	Feb 24	METADEX enhancements
NEWS	22	Feb 24	PCTGEN now available on STN
NEWS	23	Feb 24	TEMA now available on STN
NEWS	24	Feb 26	NTIS now allows simultaneous left and right truncation
NEWS	25	Feb 26	PCTFULL now contains images
NEWS	26	Mar 04	SDI PACKAGE for monthly delivery of multifile SDI results
NEWS	27	Mar 20	EVENTLINE will be removed from STN
NEWS	28	Mar 24	PATDPAFULL now available on STN
NEWS	29	Mar 24	Additional information for trade-named substances without structures available in REGISTRY
NEWS	30	Apr 11	Display formats in DGENE enhanced
NEWS	31	Apr 14	MEDLINE Reload
NEWS	32	Apr 17	Polymer searching in REGISTRY enhanced
NEWS	33	Apr 21	Indexing from 1947 to 1956 being added to records in CA/CAPLUS
NEWS	34	Apr 21	New current-awareness alert (SDI) frequency in WPIDS/WPINDEX/WPIX
NEWS	35	Apr 28	RDISCLOSURE now available on STN
NEWS	36	May 05	Pharmacokinetic information and systematic chemical names added to PHAR
NEWS	37	May 15	MEDLINE file segment of TOXCENTER reloaded
NEWS	38	May 15	Supporter information for ENCOMPPAT and ENCOMPLIT updated
NEWS	39	May 16	CHEMREACT will be removed from STN
NEWS	40	May 19	Simultaneous left and right truncation added to WSCA
NEWS	41	May 19	RAPRA enhanced with new search field, simultaneous left and right truncation

NEWS EXPRESS April 4 CURRENT WINDOWS VERSION IS V6.01a, CURRENT
 MACINTOSH VERSION IS V6.0b(ENG) AND V6.0Jb(JP),
 AND CURRENT DISCOVER FILE IS DATED 01 APRIL 2003
 NEWS HOURS STN Operating Hours Plus Help Desk Availability
 NEWS INTER General Internet Information
 NEWS LOGIN Welcome Banner and News Items
 NEWS PHONE Direct Dial and Telecommunication Network Access to STN
 NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that
 specific topic.

All use of STN is subject to the provisions of the STN Customer
 agreement. Please note that this agreement limits use to scientific
 research. Use for software development or design or implementation
 of commercial gateways or other similar uses is prohibited and may
 result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 09:36:15 ON 29 MAY 2003

=> file agricola caplus biosis

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'AGRICOLA' ENTERED AT 09:36:26 ON 29 MAY 2003

FILE 'CAPLUS' ENTERED AT 09:36:26 ON 29 MAY 2003
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'BIOSIS' ENTERED AT 09:36:26 ON 29 MAY 2003
 COPYRIGHT (C) 2003 BIOLOGICAL ABSTRACTS INC. (R)

=> s pld? or phospholipase d
 L1 11845 PLD? OR PHOSPHOLIPASE D

=> s l1 and plant?
 L2 824 L1 AND PLANT?

=> s l2 and transgenic
 L3 43 L2 AND TRANSGENIC

=> dup rem l3
 PROCESSING COMPLETED FOR L3
 L4 30 DUP REM L3 (13 DUPLICATES REMOVED)

=> d 1-10 ti

L4 ANSWER 1 OF 30 CAPLUS COPYRIGHT 2003 ACS
 TI Use of **phospholipase D** gene of Arabidopsosis thaliana
 in regulating drought tolerant of **transgenic plants**

L4 ANSWER 2 OF 30 CAPLUS COPYRIGHT 2003 ACS
 TI Protein and cDNA sequence of **plant** proteins and enzymes and
 their uses in controlling **phospholipase D** expression
 in **transgenic plants**

L4 ANSWER 3 OF 30 CAPLUS COPYRIGHT 2003 ACS
 TI Developmental regulation of **phospholipase D** in tomato
 fruits

L4 ANSWER 4 OF 30 CAPLUS COPYRIGHT 2003 ACS
 TI Expression vector construction for expression of foreign gene in chloroplasts of **plants**

L4 ANSWER 5 OF 30 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved. (2003) DUPLICATE 1
 TI T4 lysozyme and attacin genes enhance resistance of **transgenic** 'Galaxy' apple against Erwinia amylovora.

L4 ANSWER 6 OF 30 CAPLUS COPYRIGHT 2003 ACS
 TI Identification of viruses capable of breaking **transgenic** resistance of papaya conferred by the coat protein gene of Papaya ringspot virus

L4 ANSWER 7 OF 30 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
 TI A role for **PLD** beta/gamma in the N-acyl ethanolamine activation of phenylalanine-ammonia lyase gene expression.

L4 ANSWER 8 OF 30 CAPLUS COPYRIGHT 2003 ACS DUPLICATE 2
 TI Transformation of Populus tomentosa with anti-**PLD** gene

L4 ANSWER 9 OF 30 CAPLUS COPYRIGHT 2003 ACS
 TI A rapid cytokinin response assay in Arabidopsis indicates a role for **phospholipase D** in cytokinin signalling

L4 ANSWER 10 OF 30 CAPLUS COPYRIGHT 2003 ACS
 TI Antisense DNAs of nematode genes expressing in **transgenic plants** for parasite resistance

=> d ab

L4 ANSWER 1 OF 30 CAPLUS COPYRIGHT 2003 ACS
 AB Methods of growing **plants** having modified transpiration rates are provided. In particular, the present invention relates to **phospholipase D** gene of Arabidopsis thaliana in regulating drought tolerant of **transgenic plants**. Such methods permit more efficient water conservation through regulation of stomatal closure responses. Accordingly, modified **plants** can be grown in areas which were previously unsuitable for growth and **plants** which can withstand drought conditions can be grown.

=> d pi

L4	ANSWER 1 OF 30	CAPLUS	COPYRIGHT 2003 ACS	
	PATENT NO.	KIND	DATE	APPLICATION NO. DATE
	-----	---	-----	-----
PI	US 2003074692	A1	20030417	US 2001-817869 20010326

=> d in

L4 ANSWER 1 OF 30 CAPLUS COPYRIGHT 2003 ACS
 IN Wang, Xuemin; Sang, Yongming

=> d 5 ab

L4 ANSWER 5 OF 30 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States

of America. It contains copyrighted materials. All rights reserved.
(2003) DUPLICATE 1

AB Genes encoding lysozyme (T4L) from T4 bacteriophage and attacin E (attE) from *Hyalophora cecropia* were used, either singly or in combination, to construct **plant** binary vectors, **pLDB15**, **p35SAMVT4**, and **pPin2Att35SAMVT4**, respectively, for *Agrobacterium*-mediated transformation of 'Galaxy' apple, to enhance resistance to *Erwinia amylovora*. In these plasmids, the T4L gene was controlled by the cauliflower mosaic virus 35S promoter with duplicated upstream domain and the untranslated leader sequence of alfalfa mosaic virus RNA 4, and the attE gene was controlled by the potato proteinase inhibitor II (Pin2) promoter. All **transgenic** lines were screened by polymerase chain reaction (PCR) for T4L and attE genes, and a double-antibody sandwich enzyme-linked immunosorbent assay for neomycin phosphotransferase II. Amplification of T4L and attE genes was observed in reverse transcriptase-PCR, indicating that these genes were transcribed in all tested **transgenic** lines containing each gene. The attacin protein was detected in all attE **transgenic** lines. The expression of attE under the Pin2 promoter was constitutive but higher levels of expression were observed after mechanical wounding. Some T4L or attE **transgenic** lines had significant disease reduction compared to nontransgenic 'Galaxy'. However, **transgenic** lines containing both attE and T4L genes were not significantly more resistant than nontransgenic 'Galaxy', indicating that there was no in **planta** synergy between attE and T4L with respect to resistance to *E. amylovora*.

=> d d 2 ab

'D' IS NOT A VALID FORMAT

In a multifile environment, a format can only be used if it is valid in at least one of the files. Refer to file specific help messages or the STNGUIDE file for information on formats available in individual files.

REENTER DISPLAY FORMAT FOR ALL FILES (FILEDEFAULT):ab

L4 ANSWER 2 OF 30 CAPLUS COPYRIGHT 2003 ACS

AB This invention relates to protein and cDNA sequences of **phospholipase D** sequence homolog from soybean, wheat, and corn. The invention also related to protein and cDNA sequences of respiratory burst oxidase homolog RbohA, RbohB, RbohC, RbohD, RbohE, RbohF sequence homolog from soybean, wheat, rice, and corn. The invention also related to protein and cDNA sequences of tRNA 5-aminomethyl-2-thiouridylate 5'-methyltransferase sequence homolog from corn and soybean. The invention also related to protein and cDNA sequences of chromomethylase sequence homolog from Jerusalem Artichoke, corn, rice and wheat. The invention also related to protein and cDNA sequences of DNA cytosine 5-methyltransferase sequence homolog from rice, soybean, and wheat. The invention also related to protein and cDNA sequences of transcription factor TFIIF subunit .alpha. and .beta. sequence homolog from corn, rice and wheat. The invention also related to protein and cDNA sequences of asparaginyl-tRNA synthetase sequence homolog and glutamyl-tRNA synthetase sequence homolog from corn, soybean, wheat and rice. The invention also related to protein and cDNA sequences of protein EDS1 (enhanced disease susceptibility 1) sequence homolog and clathrin-assocd. AP-2 complex AP50 subunit sequence homolog from corn, rice, soybean, and wheat. The invention also related to protein and cDNA sequences of .alpha.-adaplin sequence homolog and .beta.-adaplin sequence homolog from corn, rice, soybean, and wheat. The invention also relates to the construction of a chimeric gene encoding all or a substantial portion of the **phospholipase D**, in sense or antisense orientation, wherein expression of the chimeric gene results in prodn. of altered levels of the **phospholipase D** in a transformed host cell.

=> d 2 pi

L4	ANSWER 2 OF 30	CAPLUS	COPYRIGHT 2003 ACS		
	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	US 2003003471	A1	20030102	US 2002-78770	20020219

=> d 2 in

L4 ANSWER 2 OF 30 CAPLUS COPYRIGHT 2003 ACS
IN Famodu, Omolayo O.; Miao, Guo-Hua; Simmons, Carl R.; Weng, Zude; Cahoon, Rebecca E.; Sakai, Hajime; Qun, Zhu; Thorpe, Catherine J.; Fader, Gary M.; Li, Bailin

=> d 3 ab

L4 ANSWER 3 OF 30 CAPLUS COPYRIGHT 2003 ACS
AB The catabolism of phospholipids initiated by **phospholipase D (PLD, EC 3.1.4.4)** is an inherent feature of developmental processes that include fruit growth and ripening. In cherry tomatoes (*Lycopersicon esculentum* Mill.), sol. and membrane-assocd. **PLD** activities increased during fruit development, which peaked at the mature green and orange stages. The increase in **PLD** activity was assocd. with a similar increase in the intensity of a 92 kDa band as demonstrated by western blot anal. A full-length cDNA having 2430 bp and encoding a putative polypeptide with 809 amino acids, was isolated using tomato RNA, RT-PCR and 5' and 3' rapid amplification of cloned ends (RACE). Anal. of the primary and secondary structures showed the presence of the C2 domain, the **PLD** domain and several other features characteristic of **PLD** alpha. Microtom tomato **plants** transformed with antisense **PLD** alpha cDNA, were similar to untransformed **plants** and showed normal fruit set and development. The ethylene climacteric was delayed by over 7 d in the antisense **PLD** fruits, indicative of a slower ripening process. The leaves and unripened fruits of antisense **PLD** microtom **plants** possessed lowered **PLD** activity and **PLD** protein, as demonstrated by western blotting. However, during ripening, **PLD** activity in the **transgenic** fruits was maintained at a higher level than that in the untransformed control. Immunolocalization of **PLD** in microtom tomato fruits revealed the cytosol-membrane translocation of **PLD** during fruit development. The ripe fruits of antisense **PLD** celebrity **plants** possessed lowered **PLD** expression and activity and showed increased firmness and red color. These results suggest that the expression of antisense **PLD** cDNA could be variable in different tomato varieties. The potential role of **PLD** in ethylene signal transduction events is discussed.

=> d 3 so

L4 ANSWER 3 OF 30 CAPLUS COPYRIGHT 2003 ACS
SO Plant Physiology and Biochemistry (Paris, France) (2003), 41(3), 223-240
CODEN: PPBIEX; ISSN: 0981-9428

=> d 4 ab

L4 ANSWER 4 OF 30 CAPLUS COPYRIGHT 2003 ACS
AB This invention provides a process of constructing of a expression vector for expression of foreign in chloroplasts of **plants**. The vector featured in contg. a ribosome binding site and promoter and terminator

sequences from chloroplasts. The vector can be used for foreign gene expression specifically in chloroplasts of **transgenic plants**.

=> d 8 ab

L4 ANSWER 8 OF 30 CAPLUS COPYRIGHT 2003 ACS DUPLICATE 2
AB Antisense **phospholipase D**.gamma. (**PLD**
.gamma.) gene was introduced into *Populus tomentosa* mediated by
Agrobacterium tumefaciens. The young leaves of triploid populus were used
as the material and the regeneration system of high frequency has been
established. The traditional transgene method by *Agrobacterium tumefaciens*
and obtained many **transgenic plants** of anti-
PLD.gamma. gene. Theses showed that the **transgenic**
plants can grow well on the culture medium with 0.7% NaCl.

=> d 8 so

L4 ANSWER 8 OF 30 CAPLUS COPYRIGHT 2003 ACS DUPLICATE 2
SO Yichuan (2002), 24(1), 40-44
CODEN: ICHUDW; ISSN: 0253-9772

=> d 9 ab

L4 ANSWER 9 OF 30 CAPLUS COPYRIGHT 2003 ACS
AB Seedlings of *Arabidopsis thaliana* harboring a fusion of the
cytokinin-responsive ARR5 gene promoter and the GUS reporter gene were
used for a pharmacol. approach to study cytokinin signal transduction.
The assay was shown to be rapid, sensitive, dose-dependent and highly
specific for cytokinins, both adenine and phenylurea derivs. Numerous
inhibitors of known signalling pathways were tested and some were shown to
suppress reporter gene induction. Particularly, primary alcs. that
specifically inhibit **phospholipase D (PLD)**
partially prevented cytokinin-induced GUS activity and reduced the
accumulation of ARR5 gene transcripts. This indicates a role for
PLD early during cytokinin signalling.

=> d 9 so

L4 ANSWER 9 OF 30 CAPLUS COPYRIGHT 2003 ACS
SO FEBS Letters (2002), 515(1-3), 39-43
CODEN: FEBLAL; ISSN: 0014-5793

=> d 11-20 ti

L4 ANSWER 11 OF 30 CAPLUS COPYRIGHT 2003 ACS
TI Hormone-dependent insertional mutants of *Arabidopsis thaliana* with reduced
viability and fertility

L4 ANSWER 12 OF 30 CAPLUS COPYRIGHT 2003 ACS
TI Collection of *Arabidopsis thaliana* morphological insertion mutants

L4 ANSWER 13 OF 30 AGRICOLA Compiled and distributed by the National
Agricultural Library of the Department of Agriculture of the United States
of America. It contains copyrighted materials. All rights reserved.
(2003) DUPLICATE 3

TI Involvement of a novel *Arabidopsis* **phospholipase D**,
AtPLDdelta, in dehydration-inducible accumulation of phosphatidic acid in
stress signalling.

- L4 ANSWER 14 OF 30 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
 TI Low temperatures reduce seed production in Arabidopsis **plants**.
- L4 ANSWER 15 OF 30 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved. (2003)
 TI Regulation of **plant** water loss by manipulating the expression of phospholipase Dalpha.
- L4 ANSWER 16 OF 30 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
 TI Overexpression of the Bt cry2Aa2 operon in chloroplasts leads to formation of insecticidal crystals.
- L4 ANSWER 17 OF 30 CAPLUS COPYRIGHT 2003 ACS
 TI Modification of lipid biosynthesis by DNA shuffling
- L4 ANSWER 18 OF 30 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved. (2003)
 TI Involvement of **Phospholipase D** in wound-induced accumulation of jasmonic acid in Arabidopsis.
- L4 ANSWER 19 OF 30 CAPLUS COPYRIGHT 2003 ACS DUPLICATE 4
 TI Effect of untranslated leader sequence of AMV RNA 4 and signal peptide of pathogenesis-related protein 1b on attacin gene expression, and resistance to fire blight in **transgenic** apple
- L4 ANSWER 20 OF 30 CAPLUS COPYRIGHT 2003 ACS
 TI Insertional mutagenesis in Arabidopsis thaliana: presonication of germinating seeds increases the efficiency of their transformation

=> d 13 ab

- L4 ANSWER 13 OF 30 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved. (2003) DUPLICATE 3
 AB Phospholipid metabolism is involved in **plant** responses to drought and salinity stress. To investigate the role of **phospholipase D (PLD)** and its product phosphatidic acid (PtdOH) in stress signalling, we isolated a novel **PLD** cDNA, designated AtPLDdelta, by screening a cDNA library prepared from dehydrated Arabidopsis thaliana. The AtPLDdelta protein, of 868 amino acids, has a putative catalytic domain and a C2 domain that is involved in Ca²⁺/phospholipid binding. The AtPLDdelta mRNA accumulated in response to dehydration and high salt stress. Histochemical analysis showed that the AtPLDdelta gene is strongly expressed in the vascular tissues of cotyledons and leaves under dehydration stress conditions. Under normal growth conditions, AtPLD delta was expressed in roots, leaves, stems and flowers but not in siliques. We showed that dehydration stimulates the accumulation of PtdOH. The accumulation of PtdOH in response to dehydration was significantly suppressed in AtPLD delta antisense **transgenic plants**. These results suggest that AtPLD delta may be involved in PtdOH accumulation in the dehydration stress response.

=> d 13 so

- L4 ANSWER 13 OF 30 AGRICOLA Compiled and distributed by the National

Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved.
(2003) DUPLICATE 3

SO The Plant journal : for cell and molecular biology, June 2001. Vol. 26, No. 6. p. 595-605
Publisher: Oxford : Blackwell Sciences Ltd.
ISSN: 0960-7412

=> d 14 ab

L4 ANSWER 14 OF 30 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
AB Arabidopsis thaliana (Columbia) and a **transgenic** line (anti-alpha **PLD**) with suppression of **phospholipase D** activity and improved freezing tolerance were used to examine the effects of low temperatures on their growth, specifically flowering and seed production. **Plants** were cooled at 1 C/h from 0 degreeC to -6 degreeC and held at each test temperature for 1 h and subsequently grown in a growth chamber at 22 degreeC with 10 h light to observe **plant** height, flowering time and morphology, and seed yield. All the growth characteristics measured in the wild type and anti-alpha **PLD plants** were adversely affected by exposure to low temperatures, more so in the **transgenic plants** than the wild type **plants**. The **plant** height decreased more or less linearly as temperature decreased and in the **transgenic plants** stem branching ceased after exposure to 0 degreeC or below. Flowering was delayed by low temperatures, by as much as 7 days in the **transgenic plants** exposed to -4 degreeC. Typically, the untreated **transgenic plants** produced about 8% to 10% less seeds than do their wild type counterparts, as they have fewer flowers and shorter silique. Although low temperatures reduced seed yield in both wild type and **transgenic plants**, the reduction was more severe in the **transgenic plants**. In **plants** exposed to 0 degreeC, the seed yield was reduced by 15% in the wild type and 30% in the **transgenic plants**. At -4 degreeC, seed yield was reduced by 60% in the wild type and by 90% in the **transgenic plants** at -5 degreeC. Thus, although the **transgenic plants** were freezing tolerant (killing temperature of -5.5 degreeC), they produced very little seeds below -4 degreeC. The results show the adverse effect of low temperatures, especially sub-zero temperatures, on the overall growth and seed production in Arabidopsis **plants**.

=> d 14 so

L4 ANSWER 14 OF 30 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
SO Hortscience, (June, 2001) Vol. 36, No. 3, pp. 548. print.
Meeting Info.: 98th Annual International Conference of the American Society for Horticultural Science Sacramento, California, USA July 21-25, 2001
ISSN: 0018-5345.

=> d 15 ab

L4 ANSWER 15 OF 30 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved.
(2003)
AB **Phospholipase D (PLD)** has been implicated in various processes, including signal transduction, membrane trafficking, and membrane degradation. Multiple forms of **PLD** with distinct biochemical properties have been described in the cell. In Arabidopsis,

PLDalpha and **PLDgamma**, but not **PLDbeta**, were detected in guard cells, and antisense suppression resulted in a specific loss of **PLDalpha**. The abrogation of **PLDalpha** rendered plants less sensitive to abscisic acid and impaired stomatal closure induced by water deficits. **PLDalpha**-depleted plants exhibited accelerated transpirational water loss and a decreased ability to tolerate drought stress. Overexpression of **PLDalpha** enhanced the leaf's sensitivity to abscisic acid. These findings provide molecular and physiological evidence that **PLDalpha** plays a crucial role in regulating stomatal movement and plant-water status.

=> d 15 so

- L4 ANSWER 15 OF 30 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved. (2003)
- S0 The Plant journal : for cell and molecular biology, Oct 2001. Vol. 28, No. 2. p. 135-144
Publisher: Oxford : Blackwell Sciences Ltd.
ISSN: 0960-7412

=> d 15 au

- L4 ANSWER 15 OF 30 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved. (2003)
- AU Sang, Y.; Zheng, S.; Li, W.; Huang, B.; Wang, X.

=> d 21-30 ti

- L4 ANSWER 21 OF 30 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
- TI **Phospholipase D**-mediated activation of NADPH oxidase and reactive oxygen generation in Arabidopsis.
- L4 ANSWER 22 OF 30 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
- TI Molecular cloning and functional analysis of polyphosphoinositide-dependent **phospholipase D**, **PLD**-beta, from Arabidopsis.
- L4 ANSWER 23 OF 30 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved. (2003) DUPLICATE 5
- TI Identification and characterization of a novel **plant phospholipase D** that requires polyphosphoinositides and submicromolar calcium for activity in Arabidopsis.
- L4 ANSWER 24 OF 30 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved. (2003) DUPLICATE 6
- TI Antisense suppression of **phospholipase D** alpha retards abscisic acid- and ethylene-promoted senescence of postharvest Arabidopsis leaves.
- L4 ANSWER 25 OF 30 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved.

(2003) DUPLICATE 7

TI Promoter analysis and expression of a **phospholipase D** gene from castor bean.

L4 ANSWER 26 OF 30 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved. (2003)

TI Characterization of **phospholipase D**-overexpressed and suppressed **transgenic** tobacco and Arabidopsis.

L4 ANSWER 27 OF 30 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.

TI Low temperature tolerance in **transgenic** tobacco and Arabidopsis thaliana expressing sense and antisense **phospholipase D** gene.

L4 ANSWER 28 OF 30 CAPLUS COPYRIGHT 2003 ACS

TI Cloning of cDNA for coat protein of papaya leaf-distortion mosaic potyvirus (**PLDMV**) for preparation of **PLDMV**-resistant **transgenic plants**

L4 ANSWER 29 OF 30 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.

TI Characterization of **phospholipase D**-overexpressed and suppressed **transgenic** tobacco and Arabidopsis.

L4 ANSWER 30 OF 30 CAPLUS COPYRIGHT 2003 ACS DUPLICATE 8

TI **Transgenic** "Malling 26" apple expressing the attacin E gene has increased resistance to Erwinia amylovora

=> d 21 ab

L4 ANSWER 21 OF 30 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.

=> d 21 so

L4 ANSWER 21 OF 30 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.

SO Plant Biology (Rockville), (1999) Vol. 1999, pp. 56. print.
Meeting Info.: Annual Meeting of the American Society of Plant Physiologists Baltimore, Maryland, USA July 24-28, 1999 American Society of Plant Physiologists (ASPP)

=> d 23 ag

'AG' IS NOT A VALID FORMAT

In a multifile environment, a format can only be used if it is valid in at least one of the files. Refer to file specific help messages or the STNGUIDE file for information on formats available in individual files.

REENTER DISPLAY FORMAT FOR ALL FILES (FILEDEFAULT):ab

L4 ANSWER 23 OF 30 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved. (2003) DUPLICATE 5

AB **Phospholipase D** (**PLD**; EC 3.1.4.4) has been proposed to be involved in a number of cellular processes including transmembrane signaling and membrane deterioration. **PLD** previously described from various **plant** sources generally requires millimolar ranges of calcium for optimal activity. In this study, we genetically suppressed the expression of this conventional **PLD** in Arabidopsis by introducing an antisense **PLD** cDNA. However,

both the antisense **transgenic** and wild-type **plants** showed comparable **PLD** activity in the presence of submicromolar concentrations of calcium and phosphatidylinositol 4,5-bisphosphate using phosphatidylcholine as a substrate. This new **PLD** activity was partially stimulated by phosphatidylinositol 4-phosphate, but not by other phospholipids, including phosphatidylinositol, phosphatidylserine, phosphatidylglycerol, phosphatidic acid, or phosphatidylcholine. Its requirement for polyphosphoinositides was further supported by its ability to be inhibited by neomycin. The polyphosphoinositide-dependent **PLD** requires calcium for activity, but no magnesium. The calcium stimulation was observed in the nanomolar range and reached a plateau at 5 micromolar calcium. The findings of this study demonstrate the presence of different **PLDs** that are regulated in a distinct manner in **plants**. The potential significance of a **PLD** that is regulated by polyphosphoinositides and physiological concentrations of Ca^{2+} is discussed.

=> d 23 so

- L4 ANSWER 23 OF 30 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved. (2003) DUPLICATE 5
- S0 The Journal of biological chemistry, Mar 14, 1997. Vol. 272, No. 11. p. 7048-7054
 Publisher: Bethesda, Md. : American Society for Biochemistry and Molecular Biology.
 CODEN: JBCHA3; ISSN: 0021-9258

=> d 24 a

'A' IS NOT A VALID FORMAT

In a multifile environment, a format can only be used if it is valid in at least one of the files. Refer to file specific help messages or the STNGUIDE file for information on formats available in individual files.

REENTER DISPLAY FORMAT FOR ALL FILES (FILEDEFAULT):ab

- L4 ANSWER 24 OF 30 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved. (2003) DUPLICATE 6
- AB Membrane disruption has been proposed to be a key event in **plant** senescence, and **phospholipase D** (**PLD**; EC 3.1.4.4) has been thought to play an important role in membrane deterioration. We recently cloned and biochemically characterized three different **PLDs** from Arabidopsis. In this study, we investigated the role of the most prevalent phospholipid-hydrolyzing enzyme, **PLDalpha**, in membrane degradation and senescence in Arabidopsis. The expression of **PLDalpha** was suppressed by introducing a **PLDalpha** antisense cDNA fragment into Arabidopsis. When incubated with abscisic acid and ethylene, leaves detached from the **PLDalpha**-deficient **transgenic plants** showed a slower rate of senescence than did those from wild-type and **transgenic control plants**. The retardation of senescence was demonstrated by delayed leaf yellowing, lower ion leakage, greater photosynthetic activity, and higher content of chlorophyll and phospholipids in the **PLDalpha** antisense leaves than in those of the wild type. Treatment of detached leaves with abscisic acid and ethylene stimulated **PLDalpha** expression, as indicated by increases in **PLDalpha** mRNA, protein, and activity. In the absence of abscisic acid and ethylene, however, detached leaves from the **PLDalpha**-deficient and wild-type **plants** showed a similar rate of senescence. In addition, the

suppression of **PLD** α did not alter natural **plant** growth and development. These data suggest that **PLD** α is an important mediator in phytohormone-promoted senescence in detached leaves but is not a direct promoter of natural senescence. The physiological relevance of these findings is discussed.

=> d 24 so

- L4 ANSWER 24 OF 30 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved. (2003) DUPLICATE 6
- SO The Plant cell, Dec 1997. Vol. 9, No. 12. p. 2183-2196
Publisher: [Rockville, MD : American Society of Plant Physiologists, c1989-
CODEN: PLCEEW; ISSN: 1040-4651

=> d 25 ab

- L4 ANSWER 25 OF 30 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved. (2003) DUPLICATE 7
- AB The expression of a castor bean (*Ricinus communis* L.) **phospholipase D (PLD; EC 3.1.4.4)** gene has been studied by examining its promoter activity in **transgenic** tobacco (*Nicotiana tabacum*) carrying a **PLD** promoter-glucuronidase transgene and by monitoring the levels of **PLD** mRNA in castor bean. Sequence and the 5' truncation analyses revealed that the 5' flanking region from nucleotide -1200 to -730 is required for the regulation and basal function of the **PLD** promoter. The **PLD** promoter in vegetative tissues is highly active in the rapidly growing regions such as the shoot apex and the secondary meristem producing axillary buds and vascular tissues of young leaves and stems. The **PLD** promoter activity in floral tissues was high in stigma, ovary, and pollen grains, but low in petals, sepals, the epidermis of anthers, styles, and filaments. The **PLD** promoter activity was enhanced by abscisic acid. Northern-blot analysis of **PLD** in castor bean showed that the **PLD** mRNA levels were high in young and metabolically more active tissues such as expanding leaves, hypocotyl hooks, developing seeds, and young seedlings, and they decreased in mature tissues such as fully expanded leaves and developed seeds. These patterns of expression suggest a role of **PLD** in rapid cell growth, proliferation, and reproduction.

=> d 25 so

- L4 ANSWER 25 OF 30 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved. (2003) DUPLICATE 7
- SO Plant physiology, Oct 1997. Vol. 115, No. 2. p. 387-395
Publisher: Rockville, MD : American Society of Plant Physiologists, 1926-
CODEN: PLPHAY; ISSN: 0032-0889

=> d 26 ab

- L4 ANSWER 26 OF 30 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved.

(2003)

=> d 26 so

L4 ANSWER 26 OF 30 AGRICOLA Compiled and distributed by the National
Agricultural Library of the Department of Agriculture of the United States
of America. It contains copyrighted materials. All rights reserved.
(2003)
SO [Physiology, biochemistry, and molecular biology of plant lipids], p.
345-347
Publisher: Dordrecht ; Boston : Kluwer Academic, c1997.
ISBN: 0792343794 (alk. paper).

=> d 27 ab

L4 ANSWER 27 OF 30 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.

=> d 27 spo

'SPO' IS NOT A VALID FORMAT

In a multifile environment, a format can only be used if it is valid
in at least one of the files. Refer to file specific help messages
or the STNGUIDE file for information on formats available in
individual files.

REENTER DISPLAY FORMAT FOR ALL FILES (FILEDEFAULT):so

L4 ANSWER 27 OF 30 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
SO Plant Physiology (Rockville), (1997) Vol. 114, No. 3 SUPPL., pp. 128.
Meeting Info.: PLANT BIOLOGY '97: 1997 Annual Meetings of the American
Society of Plant Physiologists and the Canadian Society of Plant
Physiologists, Japanese Society of Plant Physiologists and the Australian
Society of Plant Physiologists Vancouver, British Columbia, Canada August
2-6, 1997
ISSN: 0032-0889.

=> d 29 ab

L4 ANSWER 29 OF 30 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.

=> d 29 so

L4 ANSWER 29 OF 30 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
SO Plant Physiology (Rockville), (1996) Vol. 111, No. 2 SUPPL., pp. 133.
Meeting Info.: Annual Meeting of the American Society of Plant
Physiologists San Antonio, Texas, USA July 27-31, 1996
ISSN: 0032-0889.

=> dis his

(FILE 'HOME' ENTERED AT 09:36:15 ON 29 MAY 2003)

FILE 'AGRICOLA, CAPLUS, BIOSIS' ENTERED AT 09:36:26 ON 29 MAY 2003

L1 11845 S PLD? OR PHOSPHOLIPASE D
L2 824 S L1 AND PLANT?
L3 43 S L2 AND TRANSGENIC
L4 30 DUP REM L3 (13 DUPLICATES REMOVED)

=> s l3 and transpiration

L5 2 L3 AND TRANSPIRATION

=> dup rem l5
PROCESSING COMPLETED FOR L5
L6 2 DUP REM L5 (0 DUPLICATES REMOVED)

=> d 1-2 ti

L6 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2003 ACS
TI Use of **phospholipase D** gene of Arabidopsosis thaliana
in regulating drought tolerant of **transgenic plants**

L6 ANSWER 2 OF 2 AGRICOLA Compiled and distributed by the National
Agricultural Library of the Department of Agriculture of the United States
of America. It contains copyrighted materials. All rights reserved.
(2003)
TI Regulation of **plant** water loss by manipulating the expression of
phospholipase Dalpha.

=> d 1-2 so

L6 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2003 ACS
SO U.S. Pat. Appl. Publ., 26 pp.
CODEN: USXXCO

L6 ANSWER 2 OF 2 AGRICOLA Compiled and distributed by the National
Agricultural Library of the Department of Agriculture of the United States
of America. It contains copyrighted materials. All rights reserved.
(2003)
SO The Plant journal : for cell and molecular biology, Oct 2001. Vol. 28, No.
2. p. 135-144
Publisher: Oxford : Blackwell Sciences Ltd.
ISSN: 0960-7412

=> d so

L6 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2003 ACS
SO U.S. Pat. Appl. Publ., 26 pp.
CODEN: USXXCO

=> d pd

'PD' IS NOT A VALID FORMAT

In a multifile environment, a format can only be used if it is valid
in at least one of the files. Refer to file specific help messages
or the STNGUIDE file for information on formats available in
individual files.

REENTER DISPLAY FORMAT FOR ALL FILES (FILEDEFAULT):ti

L6 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2003 ACS
TI Use of **phospholipase D** gene of Arabidopsosis thaliana
in regulating drought tolerant of **transgenic plants**

=> s l4 and stomat?

L7 2 L4 AND STOMAT?

=> d 1-2 ti

L7 ANSWER 1 OF 2 AGRICOLA Compiled and distributed by the National
Agricultural Library of the Department of Agriculture of the United States
of America. It contains copyrighted materials. All rights reserved.
(2003)

TI Regulation of **plant** water loss by manipulating the expression of phospholipase Dalpha.

L7 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2003 ACS

TI Use of **phospholipase D** gene of Arabidopsosis thaliana in regulating drought tolerant of **transgenic plants**

=> d 1-2 so

L7 ANSWER 1 OF 2 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved. (2003)

S0 The Plant journal : for cell and molecular biology, Oct 2001. Vol. 28, No. 2. p. 135-144
Publisher: Oxford : Blackwell Sciences Ltd.
ISSN: 0960-7412

L7 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2003 ACS

S0 U.S. Pat. Appl. Publ., 26 pp.
CODEN: USXXCO

=> s l4 and abscisic acid

L8 5 L4 AND ABSCISIC ACID

=> dup rem l8

PROCESSING COMPLETED FOR L8

L9 5 DUP REM L8 (0 DUPLICATES REMOVED)

=> d 1-5 ti

L9 ANSWER 1 OF 5 CAPLUS COPYRIGHT 2003 ACS

TI Use of **phospholipase D** gene of Arabidopsosis thaliana in regulating drought tolerant of **transgenic plants**

L9 ANSWER 2 OF 5 CAPLUS COPYRIGHT 2003 ACS

TI A rapid cytokinin response assay in Arabidopsis indicates a role for **phospholipase D** in cytokinin signalling

L9 ANSWER 3 OF 5 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved. (2003)

TI Regulation of **plant** water loss by manipulating the expression of phospholipase Dalpha.

L9 ANSWER 4 OF 5 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved. (2003)

TI Antisense suppression of **phospholipase D** alpha retards **abscisic acid**- and ethylene-promoted senescence of postharvest Arabidopsis leaves.

L9 ANSWER 5 OF 5 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved. (2003)

TI Promoter analysis and expression of a **phospholipase D** gene from castor bean.

=> d 1-5 so

- L9 ANSWER 1 OF 5 CAPLUS COPYRIGHT 2003 ACS
S0 U.S. Pat. Appl. Publ., 26 pp.
CODEN: USXXCO
- L9 ANSWER 2 OF 5 CAPLUS COPYRIGHT 2003 ACS
S0 FEBS Letters (2002), 515(1-3), 39-43
CODEN: FEBLAL; ISSN: 0014-5793
- L9 ANSWER 3 OF 5 AGRICOLA Compiled and distributed by the National
Agricultural Library of the Department of Agriculture of the United States
of America. It contains copyrighted materials. All rights reserved.
(2003)
S0 The Plant journal : for cell and molecular biology, Oct 2001. Vol. 28, No.
2. p. 135-144
Publisher: Oxford : Blackwell Sciences Ltd.
ISSN: 0960-7412
- L9 ANSWER 4 OF 5 AGRICOLA Compiled and distributed by the National
Agricultural Library of the Department of Agriculture of the United States
of America. It contains copyrighted materials. All rights reserved.
(2003)
S0 The Plant cell, Dec 1997. Vol. 9, No. 12. p. 2183-2196
Publisher: [Rockville, MD : American Society of Plant Physiologists,
c1989-
CODEN: PLCEEW; ISSN: 1040-4651
- L9 ANSWER 5 OF 5 AGRICOLA Compiled and distributed by the National
Agricultural Library of the Department of Agriculture of the United States
of America. It contains copyrighted materials. All rights reserved.
(2003)
S0 Plant physiology, Oct 1997. Vol. 115, No. 2. p. 387-395
Publisher: Rockville, MD : American Society of Plant Physiologists, 1926-
CODEN: PLPHAY; ISSN: 0032-0889

=> d 2 ab

- L9 ANSWER 2 OF 5 CAPLUS COPYRIGHT 2003 ACS
AB Seedlings of Arabidopsis thaliana harboring a fusion of the
cytokinin-responsive ARR5 gene promoter and the GUS reporter gene were
used for a pharmacol. approach to study cytokinin signal transduction.
The assay was shown to be rapid, sensitive, dose-dependent and highly
specific for cytokinins, both adenine and phenylurea derivs. Numerous
inhibitors of known signalling pathways were tested and some were shown to
suppress reporter gene induction. Particularly, primary alcs. that
specifically inhibit **phospholipase D (PLD)**
partially prevented cytokinin-induced GUS activity and reduced the
accumulation of ARR5 gene transcripts. This indicates a role for
PLD early during cytokinin signalling.

=> d 4 ab

- L9 ANSWER 4 OF 5 AGRICOLA Compiled and distributed by the National
Agricultural Library of the Department of Agriculture of the United States
of America. It contains copyrighted materials. All rights reserved.
(2003)
AB Membrane disruption has been proposed to be a key event in **plant**
senescence, and **phospholipase D (PLD; EC**
3.1.4.4) has been thought to play an important role in membrane
deterioration. We recently cloned and biochemically characterized three
different **PLDs** from Arabidopsis. In this study, we investigated

the role of the most prevalent phospholipid-hydrolyzing enzyme, **PLDalpha**, in membrane degradation and senescence in Arabidopsis. The expression of **PLDalpha** was suppressed by introducing a **PLDalpha** antisense cDNA fragment into Arabidopsis. When incubated with **abscisic acid** and ethylene, leaves detached from the **PLDalpha**-deficient **transgenic plants** showed a slower rate of senescence than did those from wild-type and **transgenic control plants**. The retardation of senescence was demonstrated by delayed leaf yellowing, lower ion leakage, greater photosynthetic activity, and higher content of chlorophyll and phospholipids in the **PLDalpha** antisense leaves than in those of the wild type. Treatment of detached leaves with **abscisic acid** and ethylene stimulated **PLDalpha** expression, as indicated by increases in **PLDalpha** mRNA, protein, and activity. In the absence of **abscisic acid** and ethylene, however, detached leaves from the **PLDalpha**-deficient and wild-type **plants** showed a similar rate of senescence. In addition, the suppression of **PLDalpha** did not alter natural **plant** growth and development. These data suggest that **PLDalpha** is an important mediator in phytohormone-promoted senescence in detached leaves but is not a direct promoter of natural senescence. The physiological relevance of these findings is discussed.

=> d 5 so

- L9 ANSWER 5 OF 5 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved. (2003)
- S0 Plant physiology, Oct 1997. Vol. 115, No. 2. p. 387-395
 Publisher: Rockville, MD : American Society of Plant Physiologists, 1926-
 CODEN: PLPHAY; ISSN: 0032-0889

=> s 14 and turgidity
 L10 0 L4 AND TURGIDITY

=> s 14 and drought
 L11 3 L4 AND DROUGHT

=> d 1-3 ti

- L11 ANSWER 1 OF 3 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved. (2003)
- TI Involvement of a novel Arabidopsis **phospholipase D**, AtPLDdelta, in dehydration-inducible accumulation of phosphatidic acid in stress signalling.
- L11 ANSWER 2 OF 3 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved. (2003)
- TI Regulation of **plant** water loss by manipulating the expression of phospholipase Dalpha.
- L11 ANSWER 3 OF 3 CAPLUS COPYRIGHT 2003 ACS
- TI Use of **phospholipase D** gene of Arabidopsis thaliana in regulating **drought** tolerant of **transgenic plants**

=> d 1-3 so

- L11 ANSWER 1 OF 3 AGRICOLA Compiled and distributed by the National
Agricultural Library of the Department of Agriculture of the United States
of America. It contains copyrighted materials. All rights reserved.
(2003)
- SO The Plant journal : for cell and molecular biology, June 2001. Vol. 26,
No. 6. p. 595-605
Publisher: Oxford : Blackwell Sciences Ltd.
ISSN: 0960-7412
- L11 ANSWER 2 OF 3 AGRICOLA Compiled and distributed by the National
Agricultural Library of the Department of Agriculture of the United States
of America. It contains copyrighted materials. All rights reserved.
(2003)
- SO The Plant journal : for cell and molecular biology, Oct 2001. Vol. 28, No.
2. p. 135-144
Publisher: Oxford : Blackwell Sciences Ltd.
ISSN: 0960-7412
- L11 ANSWER 3 OF 3 CAPLUS COPYRIGHT 2003 ACS
- SO U.S. Pat. Appl. Publ., 26 pp.
CODEN: USXXCO